



CellMate[®] Foliar Plus Mo

FOLIAR-APPLIED LIQUID NUTRIENT

N 3%

Ca 8%

B 0.5%

Mo 0.2%

3-0-0

GUARANTEED ANALYSIS

Total Nitrogen (N)	3.0%
3.0% Nitrate Nitrogen	
Calcium (Ca)	8.0%
Boron (B)	0.5%
Molybdenum (Mo)	0.2%

Derived from: calcium gluconate, calcium nitrate, boric acid, and sodium molybdate.

2.5 US Gal / 9.5 Liters

11.7 lbs per Gallon @ 68° F

1.4 kgs per Liter @ 20° C

Net Weight 29.3 lbs / 13.2 kgs



DO NOT STORE BELOW 40°F (5°C)

FBSciences Inc.

153 N. Main St, Suite 100 | Collierville, TN 38017 USA

USA062422 | F1837

3102416



CellMate® Foliar Plus Mo

DIRECTIONS FOR USE

DO NOT spray in the heat of the day or when the plant is under moisture stress. DO NOT spray to the point of runoff. Use as fine of a spray mist as possible. Allow 3-4 hours after application before sprinkler irrigating to avoid washing the product off.

Recommended mixing sequence: water, adjuvants, pesticides, FBSciences nutrient products, other fertilizers, balance of water while agitating. When mixing with high phosphate fertilizers, add a citric acid buffer until the pH is 4.5 to 5.0 to improve compatibility and uptake. Ensure agitation is available when mixing with sulfate fertilizers. A standard jar test is recommended before tank mixing.

RATES & TIMING FOR ALL CROPS

Apply **1-4 quarts/acre** anytime during the growing season; use the higher rate with spray volume greater than 50 gallons per acre; repeat as needed.

INSTRUCCIONES DE USO

NO rociar en el calor del día o cuando la planta esté bajo estrés hídrico. NO rociar hasta el punto de escurrimiento. Usar una neblina de rociado tan fina como sea posible. Deje pasar de 3-4 horas después de la aplicación antes de regar con aspersión para evitar lavar el producto.

Secuencia de mezcla recomendada: agua, adyuvantes, pesticidas, productos nutritivos de FBSciences, otros fertilizantes, balance de agua mientras se agita. Al mezclar con fertilizantes altos en fosfato, agregar un buffer de ácido cítrico hasta que el pH esté de 4.5 a 5.0 para mejorar la compatibilidad y la absorción. Asegurarse de que la agitación esté disponible cuando se mezcla con fertilizantes de sulfato. Se recomienda realizar una prueba de jarra estándar antes de mezclar en el tanque.

TASAS Y MOMENTO PARA TODOS LOS CULTIVOS

Aplicar de **1 a 4 cuartos de galón/acre** en cualquier momento durante la temporada de crecimiento; usar la dosis más alta con un volumen de rociado superior a 50 galones por acre; repetir según sea necesario.

WARNING: This product contains boron. Over application or excessive use of boron, or use on crops other than those listed, can result in boron toxicity, causing foliar damage or serious plant injury.

WARNING: The application of fertilizer material containing molybdenum (Mo) may result in forage crops containing levels of molybdenum (Mo) which are toxic to ruminant animals.

WARNING

READ ENTIRE LABEL BEFORE USE | KEEP OUT OF REACH OF CHILDREN



HAZARD STATEMENTS Causes serious eye irritation.

AVISO PELIGRO Provoca irritación ocular grave.

PRECAUTIONARY STATEMENTS Wear protective gloves/clothing and eye/face protection. Wash skin thoroughly after handling. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists, get medical attention. Dispose of contents/container in accordance with local/national regulations.

WARRANTY AND CONDITIONS OF SALE Manufacturer warrants that this material conforms to the chemical description hereon and is reasonably fit for use as directed. No other warranty is made, either expressed or implied. BEFORE OPENING, see other conditions and limitations that apply which are available at: <http://www.FBSciences.com/ProductConditionsOfSale/>. See patent and other patents pending: <http://www.fbsciencespatents.com>. Information regarding the contents and levels of metals in this product is available on the Internet at: <http://www.aapfco.org/metals.html>

USA062422 | 3152416